arXiv:1609.03133v1 [math.AC] 11 Sep 2016

WHEN ALMOST COHEN-MACAULAY ALGEBRAS MAP INTO BIG COHEN-MACAULAY MODULES

RAJSEKHAR BHATTACHARYYA

Abstract. In this paper, we show that almost Cohen-Macaulay algebras are solid and in this respect, we seek for some special situation when (a) an almost Cohen-Macaulay algebra will be phantom extension and (b) when it maps into a balanced big Cohen-Macaulay module.

1. introduction

Let $(R, \mathfrak{m})$ be a Noetherian local ring. An R-algebra B is defined as big Cohen-Macaulay algebra if some system of parameters of R is a regular sequence on B. It is balanced if every system of parameters of R is a regular sequence on B. In equal characteristic, the tight closure operation has been used to present proofs of the existence of balanced big Cohen-Macaulay modules and algebras.

In [Di1], a list of seven axioms, which are also known as "finite axioms" for a closure operation, is defined for finitely generated modules over a complete local domain R. In a recent work of Dietz [Di2], these axioms are extended beyond finitely generated modules and we call them as "big axioms". Any closure operation which satisfies these axioms is called Dietz closure. A Dietz closure is powerful enough to produce big Cohen-Macaulay modules.

We recall the definition of solid algebra and module (see [Ho]) and definitions phanton extension for a closure operation (see [HH3] and [Di1]). In [Di2], the notion of phantom extension is extended for closure operation which also satisfies "big axioms" and there it is proved that every phantom extension is solid, but converse is not true.

Let R be a Noetherian local domain and R^+ be its integral closure in an algebraic closure of its fraction field. We observe the definition of an almost Cohen-Macaulay algebra R^+-algebra [Ro] (see Section 2). It is a well-known result due to Hochster that for a complete local domain, any algebra which maps into big Cohen-Macaulay algebra is solid. In this paper, we show that almost Cohen-Macaulay algebras are solid. Moreover, here, we seek for some special situation when (a) an almost Cohen-Macaulay algebra will be phantom extension and (b) when it maps into a balanced big Cohen-Macaulay module. More precisely, in this paper we show that:

(1) Over a complete local domain R of mixed characteristic $p > 0$, an almost Cohen-Mcaulay R^+-algebra S, is a solid module. As an immediate consequence of this result, we get that for an almost Cohen-Macaulay algebra S over a complete local domain R with F-finite residue field, there always exists a big Cohen-Macaulay module B' over some complete local domain R', such that R' is homomorphic image of R and S is mapped into B' (see Theorem 3.1 and Corollary 2).

(2) Over a complete local domain R of mixed characteristic $p > 0$, an almost Cohen-Mcaulay R^+-algebra S (with certain restrictions on S) is a phantom extension of R via the closure defined in Definition 2 (see Section 2) and moreover, we observe that S can be modified into a balanced big Cohen-Macaulay module over R (see Theorem 4.1 and Corollary 3).

2. PRELIMINARY RESULTS

Let R be a Noetherian local domain of $\dim d$ and R^+ be its integral closure in an algebraic closure of its fraction field. Let R^+ be equipped with a normalized value map. Consider a local R^+-algebra A. We recall the following definition of a closure operation from first paragraph of Section 5 of [AB], which extend the notion of dagger closure in [HH1], via a local algebra over R^+.

Definition 1. Let A be a local R^+-algebra where R^+ is equipped with a normalized valuation $v : R^+ \to \mathbb{R} \cup \{\infty\}$ and M be an A-module. Consider a submodule $N \subset M$. We say whenever $x \in N_M^v$ if and only if for every $\epsilon > 0$, there exists $a \in R^+$ such that $v(a) < \epsilon$ and $ax \in N$ or sometimes we can say there exists $a \in R^+$ with arbitrarily small $v(a)$ and $ax \in N$. A submodule N is called closed if $N = N_M^v$.

Throughout this work we fix v. We recall the definition of almost Cohen-Macaulay R^+-algebra [Ro]: Let R^+ is equipped with a value map. An R^+-algebra is almost Cohen-Macaulay if for every system of parameters $x_1, \dots, x_d$, $((x_1, \dots, x_i) : x_{i+1})/(x_1, \dots, x_i)$ is almost zero R^+-module for every $i = 1 \to d-1$ and $A/\mathfrak{m}A$ is not almost zero. If we do not put any restriction on $A/\mathfrak{m}A$, we call it an weakly almost Cohen-Macaulay [Sh].

We list the following properties of the above closure. The proof of all these properties are given in [AB], here we restate it for completeness.

Proposition 2.1. *Let M, M' be modules over a local R^+-algebra A with a normalized valuation $v : R^+ \to \mathbb{R} \cup \{\infty\}$. Consider the arbitrary submodules N, W of M. Then the following are true:*

(i) *N_M^v is a submodule of M containing N.*
(ii) *$(N_M^v)_M^v = N_M^v$.*
(iii) *If $N \subset W \subset M$, then $N_M^v \subset W_M^v$.*
(iv) *Let $f : M \to M'$ be a homomorphism. Then $f(N_M^v) \subset f(N)_{M'}^v$.*
(v) *If $N_M^v = N$ then $0_{M/N}^v = 0$.*

In addition to, if A is weakly almost Cohen-Macaulay then following is true:

(vi) *Let $x_1, \dots, x_{k+1}$ be a partial system of parameters for A, and let $J = (x_1, \dots, x_k)A$. Suppose that there exists a surjective homomorphism $f : M \to A/J$ such that $f(u) = \bar{x}_{k+1}$, where $\bar{x}$ is the image of x in A/J. Then $(Au)_M^v \cap ker f \subset (Ju)_M^v$.*

Proof. (i) Clearly $N \subset N_M^v$. For $x, y \in N_M^v$, take $\epsilon > 0$ and choose $a, b \in R^+$ such that $v(a), v(b) < \epsilon/2$ and $ax, by \in N$. Thus we have $v(ab) < \epsilon$ and $ab(x+y) \in N$. Thus $x+y \in N_M^v$. Consider $x \in N_M^v$ and $b \in A$. Since there exists $a \in R^+$ such that $v(a) < \epsilon$ and $ax \in N$ and since N is a submodule, we find $a(bx) \in N$ and $bx \in N_M^v$. Thus it is easy to see that N_M^v is a submodule containing N.

(ii) Take $x \in (N_M^v)_M^v$. For $\epsilon > 0$ and choose $a \in R^+$ such that $v(a) < \epsilon/2$ and $ax \in (N_M^v)$. Similarly, for $\epsilon > 0$ and choose $b \in R^+$ such that $v(b) < \epsilon/2$ and $(ba)x \in N$. Thus we find $ba \in R^+$ with $v(ba) < \epsilon$ such that $(ba)x \in N$. So $x \in (N_M^v)$, which yields the claim.

(iii) This is easy and we leave it to reader.

(iv) Consider $x \in N_M^v$, thus for every $\epsilon > 0$, there exists $a \in R^+$ such that $v(a) < \epsilon$ and $ax \in N$. This implies $f(ax) = af(x)$ is in $f(N)$ where $a \in R^+$ is of arbitrarily small positive order. So $f(x) \in f(N)_{M'}^v$.

(v) Consider $\bar{x} \in 0_{M/N}^v$ which is the image of x in M/N. This implies that $ax \in N$ for the element $a \in R^+$ of arbitrarily small positive order. So $x \in N_M^v = N$ and $\bar{x} \in 0$.

(vi) Take $x \in (Au)_M^v \cap \ker f$. For every $\epsilon > 0$ there exists $a \in R^+$ of $v(a) < \epsilon/2$ such that $ax = bu \in Au$ and $af(x) = 0 = bf(u) = b\bar{x}_{k+1}$. This implies $bx_{k+1} \in J$ i.e. $b \in (J : x_{k+1})$. Since A is almost Cohen-Macaulay, for every $\epsilon > 0$ there exists $c \in R^+$ of $v(c) < \epsilon/2$ such that $cb \in J$. Thus for $\epsilon > 0$ there exists $ac \in R^+$ of $v(ac) < \epsilon$ and $(ac)x = cbu \in Ju$. So $x \in (Ju)_M^v$.

□

With help of above closure, we can define a closure operation for submodules of an arbitrary R-module M. This definition is similar to that, given in the first paragraph of Section 5 of [AB], where it was given for finitely generated R modules.

Definition 2. Let $(R, \mathfrak{m})$ be a Noetherian local domain and let A be a local R^+-algebra and R^+ is equipped with a normalized valuation $v : R^+ \to \mathbb{R} \cup \{\infty\}$. For any R-module M and for its submodule N we define submodule $N_M^{\mathbf{v}}$ such that $x \in N_M^{\mathbf{v}}$ if $x \otimes 1 \in \operatorname{im}(A \otimes N \to A \otimes M)_{A\otimes M}^v$.

Remark 1. *The definition of above closure depends not only on the value map but also on the R^+-algebra A. Here, we fix this notation i.e. closure operation will always be performed by A.*

We recall the definition of Dietz closure [Di1], and also big Axioms of Dietz, see definition 1.1 of [Di2]. We observe the following proposition.

Proposition 2.2. *Let $(R, \mathfrak{m})$ be a Noetherian local domain and let R^+ be equipped with a normalized valuation $v : R^+ \to \mathbb{R} \cup \{\infty\}$. Let M, M' be arbitrary R-modules. Consider the submodules N, W of M. Then the following are true:*

(i) *$N_M^{\mathbf{v}}$ is a submodule of M containing N.*

(ii) *$(N_M^{\mathbf{v}})_M^{\mathbf{v}} = N_M^{\mathbf{v}}$.*

(iii) *If $N \subset W \subset M$, then $N_M^{\mathbf{v}} \subset W_M^{\mathbf{v}}$.*

(iv) *Let $f : M \to M'$ be a homomorphism. Then $f(N_M^{\mathbf{v}}) \subset f(N)_{M'}^{\mathbf{v}}$.*

(v) *If $N_M^{\mathbf{v}} = N$ then $0_{M/N}^{\mathbf{v}} = 0$.*

(vi) *If $R^+ \to A$ is injective and $A/\mathfrak{m}A$ is not almost zero then $0^{\mathbf{v}} = 0$ ideal and $m^{\mathbf{v}} = m$. In particular closure of every $I \subset R$ is proper.*

In addition to, if A is weakly almost Cohen-Macaulay then following is true:

(vii) *Let $x_1, \ldots, x_{k+1}$ be a partial system of parameters for R, and let $J = (x_1, \ldots, x_k)R$. Suppose that there exists a surjective homomorphism $f : M \to R/J$ such that $f(u) = \bar{x}_{k+1}$, where $\bar{x}$ is the image of x in R/J. Then $(Ru)_M^{\mathbf{v}} \cap ker f \subset (Ju)_M^{\mathbf{v}}$.*

In other words, if A is an almost Cohen-Macaulay R^+-algebra where R^+ is contained in it as a subdomain, then above closure operation satisfies big Axioms of Dietz.

Proof. (i) Clearly $N \subset N_M^{\mathbf{v}}$. To prove $N_M^{\mathbf{v}}$ is a submodule, let $x, y \in N_M^{\mathbf{v}}$ and $r \in R$. Then

$$x \otimes 1, y \otimes 1 \in \operatorname{im}(A \otimes N \to A \otimes M)^v_{A \otimes M}.$$

Thus $(x+y) \otimes 1, rx \otimes 1 \in \operatorname{im}(A \otimes N \to A \otimes M)^v_{A \otimes M}$. These yield that $x + y, rx \in N_M^{\mathbf{v}}$.

(ii) Let $x \in (N_M^{\mathbf{v}})_M^{\mathbf{v}}$. This implies $x \otimes 1 \in \operatorname{im}(A \otimes N_M^{\mathbf{v}} \to A \otimes M)^v_{A \otimes M}$. Equivalently, for $\epsilon > 0$ there exists $a \in R^+$ such that $v(a) < \epsilon/2$ and $a \otimes x = \sum_{i=1}^{l} a_i \otimes x_i$ where $x_i \in N_M^{\mathbf{v}}$ and $a_i \in A$. Choose $c_i \in R^+$ such that for every i, $v(c_i) < \epsilon/2l$ and $x_i \otimes c_i \in \operatorname{im}(A \otimes N \to A \otimes M)$. Take $c = \prod_{i=1}^{l} c_i$. Thus $x \otimes ac \in \operatorname{im}(A \otimes N \to A \otimes M)$ and $x \in (N_M^{\mathbf{v}})$.

(iii) This is trivial.

(iv) Let $x \in N_M^{\mathbf{v}}$ and this gives $a \otimes x = \sum_{i=1}^{l} a_i \otimes x_i$ for element $a \in R^+$ of arbitrarily small order, where $a_i \in A$ and $x_i \in N$. Applying $f \otimes 1_A$, we get $a \otimes f(x) = \sum_{i=1}^{l} a_i \otimes f(x_i)$ and thus we conclude.

(v) Denote the image of $x \in M$ in M/N by $\bar{x}$. If $\bar{x} \in 0^{\mathbf{v}}_{M/N}$ then $a \otimes \bar{x} \in 0$ and this implies $a \otimes x \in \operatorname{im}(A \otimes N \to A \otimes M)$ for element $a \in R^+$ of arbitrarily small order. Thus $x \in N_M^{\mathbf{v}} = N$ and $\bar{x} \in 0$.

(vi) Due to (e) of Lemma 1.3 of [Di2], the condition $0^{\mathbf{v}} = 0$ follows. For $m^{\mathbf{v}} = m$, suppose on the contrary that $s \in \mathfrak{m}^{\mathbf{v}}$ for some unit element s. It turns out that $sc \in \mathfrak{m}A$ for every element c of arbitrarily small order, i.e., $\mathfrak{m}A$ contain elements of arbitrarily small order, since s is a unit. This provides a contradiction, because A/mA is not almost zero.

(vii) Take $x \in (Ru)_M^{\mathbf{v}} \cap \ker f$. For every $\epsilon > 0$ there exists $a \in R^+$ of $v(a) < \epsilon/2$ such that $a \otimes x = \sum_{i=1}^{n} a_i \otimes r_i u$. Since $f(x) = 0$, $f(x) \otimes a = x_{k+1}(\sum_{i=1}^{n} r_i a_i) + JA = 0$. This implies $\sum_{i=1}^{n} r_i a_i \in (JA : x_{k+1}A) \subset (JA)^v$. So there exists $c \in R^+$ with $v(c) < \epsilon/2$ such that $c \sum_{i=1}^{n} r_i a_i \in JA$. Thus

$$ac \otimes x = u \sum_{i=1}^{n} r_i a_i \otimes c \in \operatorname{im}(JA \otimes Ru \to A \otimes M)$$

and this gives $ac \otimes x \in \operatorname{im}(A \otimes Ju \to A \otimes M)$ with $v(ac) < \epsilon$. So we conclude.

□

As a consequence of above proposition we have the following corollary where we observe that the condition $0^{\mathbf{v}} = 0$ is equivalent to the condition that R^+ should be inside of A as a subdomain.

Corollary 1. *Let (R, m) be a Noetherian local domain and let R^+ be equipped with a normalized valuation $v : R^+ \to \mathbb{R} \cup \{\infty\}$. Consider a local R^+-algebra A and the closure operation defined as above. Under this situation, closure operation satisfies all the properties given in Proposition 2.2 if and only if A is an almost Cohen-Macaulay R^+-algebra where R^+ is contained in it as a subdomain.*

Proof. From Proposition 2.2 'if' follows. Here we only prove 'only if'. In above situation it is clear that $m^{\mathbf{v}} = m$ implies A/mA is not almost zero. Moreover $0^{\mathbf{v}} = 0$ also implies $R^+ \to A$ is injective: take $0 \neq a \in R^+$ such that its image in A is zero. Since a is integral over R, let there exists a minimal monic expression $a^n + r_1 a^{n-1} + \ldots + r_n = 0$, where each $r_i \in R$ with $r_n \neq 0$. Take the image of the expression in A which gives image of r_n is zero in A. So $r_n \in 0^{\mathbf{v}} = 0$. This gives $r_n = 0$. So we arrive at a contradiction and a is zero in R^+. Moreover, since this closure captures colon in A, we can conclude. □

3. almost cohen-macaulay algebras are solid

We recall the definition of solid module and algebra [Ho]. It is a well-known result due to Hochster (see, Corollary 10.6 of [Ho]) that for a complete local domain, any algebra which maps into big Cohen-Macaulay algebra is solid. Similar is true for almost Cohen-Macaulay algebra.

Theorem 3.1. *(a) Let R be a complete local domain and S be an almost Cohen-Macaulay algebra. Then S is a solid R-algebra.*
(b) In particular, any algebra which maps into an almost Cohen-Macaulay algebra is solid.
(c) Let $R^+ = \lim_{\to} R_\alpha$ be a direct limit of Noetherian local rings R_α, where each R_α is module finite over R. If S is almost Cohen-Macaulay algebra, then S is a solid R_α-algebra for each α.

Proof. (a) Let $(R, \mathfrak{m})$ be a complete local domain of dimension d and S be an almost Cohen-Macaulay algebra. Let $H^d_{\mathfrak{m}S}(S)$ be the local cohomology module in the support of $\mathfrak{m}$ and we can view it as $\lim_{\to} S/(x_1^t, \ldots, x_d^t)S$ with multiplication map "$x_1 \ldots x_d$". We would like to show $\lim_{\to} S/(x_1^t, \ldots, x_d^t)S \neq 0$. From definition $S/(x_1, \ldots, x_d)S$ is nonzero. If $\lim_{\to} S/(x_1^t, \ldots, x_d^t)S = 0$, then image of 1 in some $S/(x_1^{t+1}, \ldots, x_d^{t+1})S$ will be zero. This implies $x_1^t \ldots x_d^t \in (x_1^{t+1}, \ldots, x_d^{t+1})S$. Set $x_1^t \ldots x_d^t = s_1 x_1^{t+1} + \ldots + s_d x_d^{t+1}$. This gives $x_1^t(x_2^t \ldots x_d^t - s_1 x_1) \in (x_2^{t+1}, \ldots, x_d^{t+1})S$. Since S is almost Cohen-Macaulay $x_1^t, x_2^{t+1} \ldots, x_d^{t+1}$ are almost regular sequence, for $\epsilon/d > 0$ there exists $b_1 \in R^+$ such that $v(b_1) < \epsilon/d$ and $b_1(x_2^t \ldots x_d^t - s_1 x_1) \in (x_2^{t+1}, \ldots, x_d^{t+1})S$ which yields $b_1(x_2^t \ldots x_d^t) \in (x_1, x_2^{t+1}, \ldots, x_d^{t+1})S$. We now repeat the process for x_2 and in a similar way for $\epsilon/d > 0$ we can get $b_2 \in R^+$ such that $v(b_2) < \epsilon/d$ and $b_1 b_2(x_3^t \ldots x_d^t) \in (x_1, x_2, x_3^{t+1}, \ldots, x_d^{t+1})S$. Thus repeating this process for d times we get $b = b_1 \ldots b_d \in (x_1, \ldots, x_d)S$ with $v(d) < \epsilon$. Thus, we get that $(x_1, \ldots, x_d)S$ as well as $\mathfrak{m}S$ contains elements of arbitrarily small order. This implies that $S/\mathfrak{m}S$ is almost zero. This is a contradiction and thus we prove $H^d_{\mathfrak{m}S}(S)$ is non-zero. Now using Corollary 2.4 of [Ho] we can conclude that S is a solid R-algebra.

(b) This is straight forward.

(c) Since for each index α, maximal ideal $\mathfrak{m}_\alpha$ and $\mathfrak{m}R_\alpha$ are m_α-primary, $H^d_{\mathfrak{m}S}(S) = H^d_{\mathfrak{m}_\alpha S}(S)$ and the assertion follows. □

4. almost cohen-macaulay algebras map into big cohen-macaulay algebra

In this section, we study when an almost Cohen-Macaulay algebra maps to big Cohen-Macaulay module. We begin the section with the following corollary which is a consequence of Theorem 3.1 of the previous section.

Corollary 2. *Let R be a complete local domain of mixed characteristic $p > 0$ with F-finite residue field and let S be an almost Cohen-Macaulay R^+-algebra. Then there exists a local domain R' (more precisely, R' turns out to be R/pR), such that S/pS is mapped into some balanced big Cohen-Macaulay module B' over R'.*
More precisely, for an almost Cohen-Macaulay R^+-algebra S over a complete local domain R with F-finite residue field, there always exists a big Cohen-Macaulay module B' over some complete local domain R', such that R' is homomorphic image of R and S is mapped into B'.

Proof. From Theorem 3.1, we have that S is solid. Passing to the ring R/pR, we have S/pS is solid as R/pR-algebra, which can be also viewed as $(R/pR)^+$-algebra. Being complete, R/pR is F-finite, since from hypothesis, residue field of R/pR which is also the residue field of R is F-finite. Now using Corollary 4.9 of [Di2] S/pS can be modified through a sequence of maps into a balanced big Cohen-Macaulay R/pR-module. Thus we conclude.

Second assertion is immediate. □

Remark 2. *For positive integers i, consider the sets $\{\epsilon_i : \epsilon_i \in \mathbb{R}\}$ and $\{c_i : c_i \in R^+\}$ such that $\lim_{i\to\infty} \epsilon_i = 0$ and $v(c_i) < \epsilon_i$. Let $R^+ = \lim_{\to} R_\alpha$ be a direct limit of Noetherian local rings R_α, where each R_α is module finite over R. If $c_i \in R_\alpha$ for some α, then we call that R_α as R_i and it is always possible to choose R_i's from the directed set of Noetherian local rings such that for $i < j$, $R_i \subset R_j$. From above Proposition 3.1, it is clear that for every positive integer i, almost Cohen-Macaulay algebra S is a solid R_i algebra.*

We propose a special kind of almost Cohen-Macaulay algebra.

Definition 3. Consider the situation of Remark 2 above. We call an almost Cohen-Maculay R^+-algebra S satisfies (*) if for every positive integer i, there exists an R_i-homomorphism h_i from S to R_i, such that it sends 1 to c_i.

Clearly, each above h_i is also R-homomorphism.

Theorem 4.1. *Let $(R, \mathfrak{m})$ be a local domain, and let S be an almost Cohen-Macaulay algebra satisfies (*). Then then S is a phantom extension of R via the closure defined in Definition 2.*

Proof. From previous Proposition S is also solid R-algebra. Since R is a domain, R/I can not be a solid algebra for any proper ideal I and so $\beta : R \to S$ must be an injection. So, we have a short exact sequence $0 \to R \to S \to Q \to 0$. Consider the following diagram, where bottom row is the projective (equivalently free) resolution of Q:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & R & \xrightarrow{\beta} & S & \longrightarrow & Q & \longrightarrow & 0 \\ & & \uparrow{\scriptstyle f} & & \uparrow{\scriptstyle g} & & \uparrow{\scriptstyle I} & & \\ \ldots & \longrightarrow & G & \xrightarrow{\nu} & F & \longrightarrow & Q & \longrightarrow & 0 \end{array}$$

Consider the sets $\{\epsilon_i : \epsilon_i \in \mathbb{R}\}$ and $\{c_i : c_i \in R^+\}$ such that $\lim_{i\to\infty} \epsilon_i = 0$ and $v(c_i) < \epsilon_i$. Now for every integer n, consider R algebra R_n, which is module finite over R and tensoring with last short exact sequence, again we get a short exact sequence $0 \to R_n \to R_n \otimes_R S \to R_n \otimes_R Q \to 0$, since by Proposition 2.1 (e) of [Ho], $R_n \otimes_R S$ is solid over R_n and for any proper ideal $J \subset R_n$, R_n/J is not solid.

From Remark 2 above, we get that for every positive integer n, S is a solid R_n-algebra, with an R_n-linear map $h_n : S \to R_n$. This h_n is also an R-linear map, which sends $h_n(1) = c_n \neq 0$, where $v(c_n) < \epsilon_n$. Thus, we can define an R-linear map $\gamma_n : R_n \otimes S \to R_n$ given by $\gamma_n(r' \otimes s) = r'h_n(s)$. Clearly, $\gamma_n(r' \otimes 1) = r'c_n$

So, for every $\epsilon > 0$, there exists an element $c_n \in R^+$, such that $v(c_n) < \epsilon_n < \epsilon$ and we have the following diagram where

$$\begin{array}{ccccccccc} 0 & \longrightarrow & R_n & \xrightarrow{1\otimes\beta} & R_n \otimes_R S & \longrightarrow & R_n \otimes_R Q & \longrightarrow & 0 \\ & & {\scriptstyle 1\otimes f}\uparrow & & \uparrow{\scriptstyle 1\otimes g} & & \uparrow{\scriptstyle I} & & \\ \ldots & \longrightarrow & R_n \otimes_R G & \xrightarrow{1\otimes\nu} & R_n \otimes_R F & \longrightarrow & R_n \otimes_R Q & \longrightarrow & 0 \end{array}$$

Set $\mathrm{Hom}_R(F, R) = F^V$ and $Im(\mathrm{Hom}_R(F, R) \to \mathrm{Hom}_R(G, R)) = Im(\nu^V)$. Now, $c_n(1 \otimes f) = \gamma_n \circ (1 \otimes \beta) \circ (1 \otimes f) = \gamma_n \circ (1 \otimes g) \circ (1 \otimes \nu) = (1 \otimes \nu)^V(\gamma_n \circ (1 \otimes g))$ and so $c_n(1 \otimes f) \in Im((1 \otimes \nu)^V)$ for all n. In other words, $c_n(1 \otimes f) \in Im(\mathrm{Hom}_{R_n}(R_n \otimes_R F, R_n) \to \mathrm{Hom}_{R_n}(R_n \otimes_R G, R_n))$

By Lemma 4.3 of [Di2] we get that $c_n(1 \otimes f) \in Im(R_n \otimes_R \mathrm{Hom}_R(F, R) \to R_n \otimes_R \mathrm{Hom}_R(G, R))$. This implies for each n, $c_n(1 \otimes f) \in Im(A \otimes_R \mathrm{Hom}_R(F, R) \to A \otimes_R \mathrm{Hom}_R(G, R))$ via base change, since A is an R_n-algebra. From above, we get for each n, $v(c_n) < \epsilon_n$, and that the ϵ_n form a sequence of real numbers converging to zero. Thus, for all $\epsilon > 0$, there exists an n such that $v(c_n) < \epsilon$ with $c_n(1 \otimes f) \in Im(A \otimes_R \mathrm{Hom}_R(F, R) \to A \otimes_R \mathrm{Hom}_R(G, R))$. Hence from Definition 2, $f \in (Im(\nu^V))^{\mathfrak{v}}_{G^V}$. From Proposition 2.2, we get that the almost Cohen-Macaulay algebra S is a phantom extension of R via closure operation defined in Definition 2. We therefore conclude. □

Corollary 3. *Let $(R, \mathfrak{m})$ be a complete local domain of mixed characteristic $p > 0$. Let S be an almost Cohen-Macaulay R-algebra which satisfies (*) condition, then S can be modified into a balanced big Cohen-Macaulay module over R.*

Proof. Using Theorem 3.7 of [Di2], Proposition 2.2 and Theorem 4.2 above, we can conclude. □

Acknowledgement . I would like thank Geoffrey D. Dietz for careful reading of the paper and for all invaluable comments and suggestions for improvement of the paper.

References

[AB] M. Asgharzadeh and R. Bhattacharyya, *Applications of closure operations on big Cohen-Macaulay algebras*, J. Algebra Appl. 11 (2012), no. 4.

[Di1] G. Dietz, *Characterization of closure operation that induced big Cohen-Macaulay algebra*, Proc. AMS **138** (2010), 3849-3862.

[Di2] G. Dietz, *Axiomatic closure operations, Phantom extensions, and Solidity*, arXiv:1511.04286.

[Ho] M. Hochster, *Solid Closure* In: Commutative algebra: syzygies, multiplicities, and birational algebra, Contemp. Math., vol. 159, pp. 103-172.

[HH1] M Hochster and C. Huneke, *Tight closure and elements of small order in integral extensions**, Journal of Pure and Applied Algebra 71 (1991) 233-247.

[HH2] M Hochster and C. Huneke, *Infinite integral extensions an big Cohen-Macaulay algebras*, Annals of Math. 135 (1992), 53-89.

[HH3] M. Hochster and C. Huneke, *Tight closure of parameter ideals and splitting in module finite extensions*, J. Algebraic Geom. 3 (1994), no. 4, 599-670

[Ro] P. Roberts, *Fontaine rings and local cohomology*, Journal of Algebra 323 (2010), 2257-2269.

[Sh] K. Shimomoto, *Almost Cohen-Macaulay algebras in mixed characteristic via Fontaine rings*, Volume 55, Number 1 (2011), 107-125, Illinois J. Math.

Dinabandhu Andrews College, Garia, Kolkata 700084, India
E-mail address: rbhattacharyya@gmail.com